


Summary of the 2012 ERA-EDTA Registry Annual Report



National and regional renal registries that contributed data to the 2012 ERA-EDTA Registry Annual Report





Renal registries contributing with individual patient data

Renal registries contributing with aggregated data

Incident patients accepted for RRT in 2012, at day 1 by country






by country




Unadjusted incidence, by country

renal registries providing individual patient data



Unadjusted incidence, by country



by country, adjusted for age and gender




Adjusted incidence, by country

renal registries providing individual patient data



Adjusted incidence, by country



mean age




Mean age at start RRT

renal registries providing individual patient data



Mean age at start RRT



mean age by gender




Mean age at start RRT

male patients starting RRT in 2012

Romania Bosnia and Herzegovina Serbia United Kingdom, Scotland Spain, Region of Murcia 61.1 Finland Spain, Andalusia Spain, Cantabria United Kingdom, England Iceland Spain, Basque country Sweden United Kingdom, Northern Ireland Croatia United Kingdom, Wales Spain, Extremadura Denmark Spain, Catalonia 64.0 Spain, Community of Madrid Spain, Valencian region Austria 64.3 Norway 64.3 The Netherlands Spain, Castile-La Mancha 64.6 Slovenia Spain, Asturias Spain, Galicia Spain, Aragon Belgium, French-speaking Spain, Castile and León France Greece Belgium, Dutch-speaking All countries 55 70 75 60 Mean age (years)

Mean age at start RRT

female patients starting RRT in 2012




by cause of renal failure




Incidence, by cause of renal failure

all patients starting RRT in 2012



Incidence, by cause of renal failure

all patients starting RRT in 2012



by cause of renal failure and age category




Incidence, by cause of renal failure

patients younger than 65 years of age at start RRT in 2012

Missing, 2.5% Glomerulonephritis/ sclerosis, 16.3% Unknown, 15.4% Pyelonephritis, 5.8% Miscellaneous, Polycystic kidneys, 18.2% adult type, 9.1% Diabetes mellitus. 22.5% Renal vascular disease, 1.4% Hypertension, 8.8%

Incidence, by cause of renal failure

patients older than 65 years of age at start RRT in 2012




by established modality




Incidence at day 91, by established modality

all patients starting RRT in 2012



Incidence at day 91, by established modality

all patients starting RRT in 2012



by established modality and age category




Incidence at day 91, by established modality

patients younger than 65 years of age at start RRT in 2012

Peritoneal dialysis, 16.9% Haemodialysis, 72.4%

Incidence at day 91, by established modality

patients older than 65 years of age at start RRT in 2012




by country




Unadjusted prevalence, by country

renal registries providing individual patient data



Unadjusted prevalence, by country




by country, adjusted for age and gender




Adjusted prevalence, by country

renal registries providing individual patient data



Adjusted prevalence, by country



mean age at December 31




Mean age at start RRT

renal registries providing individual patient data



Mean age at start RRT



mean age by gender at December 31




Mean age at start RRT

male patients starting RRT in 2012

United Kingdom, Scotland 56.4 Spain, Community of Madrid 56.6 Spain, Valencian region **57.4** Belgium, Dutch-speaking 57.6 Spain, Region of Murcia 57.8 United Kingdom, Northern Ireland; 58.2 Croatia 58.3 United Kingdom, England 58.3 Denmark 58.7 Spain, Cantabria 58.7 Spain, Galicia 59.0 United Kingdom, Wales 59.1 Slovenia 59.3 Spain, Extremadura Sweden Spain, Catalonia 60.1 Austria 60.2 Spain, Aragon The Netherlands Greece Romania Norway Spain, Asturias Iceland Spain, Andalusia Spain, Castile and León Spain, Basque country Finland France Spain, Castile-La Mancha 62.9 Serbia Bosnia and Herzegovina Belgium, French-speaking All countries 60.2 55 60 65 70 Mean age

Mean age at start RRT

female patients starting RRT in 2012




by cause of renal failure




Prevalence, by cause of renal failure

all patients on RRT in 2012



Prevalence, by cause of renal failure

all patients on RRT in 2012




by cause of renal failure and age category




Prevalence, by cause of renal failure

patients younger than 65 years of age in 2012



Prevalence, by cause of renal failure

patients older than 65 years of age in 2012




by established modality




Prevalence, by established modality

all patients on RRT in 2012



Prevalence, by established modality

all patients on RRT in 2012




by established modality and age category




Prevalence, by established modality

patients younger than 65 years in 2012



Prevalence, by established modality

patients older than 65 years of age in 2012




Renal transplants performed in 2012

by donor type




Transplantation activity, by donor type

all patients transplanted in 2012



Transplantation activity, by donor type

all patients transplanted in 2012



Renal transplants performed in 2012

by donor type and age category




Transplantation activity, by donor type

patients younger than 65 years of age at transplantation

Living donor, 23.5% Deceased donor, 71.5%

Transplantation activity, by donor type

patients older than 65 years of age at transplantation



Survival probability


by modality



Adjusted survival (cohort 2003-2007)

incident dialysis patients and patients receiving a first transplant

from day 91, by modality adjusted for age, gender and cause of renal failure



Survival probability


by modality and primary diagnosis



Adjusted survival (cohort 2003-2007)

incident haemodialysis patients

from day 91, by modality adjusted for age, gender and cause of renal failure



Adjusted survival (cohort 2003-2007)

incident peritoneal dialysis patients

from day 91, by modality adjusted for age, gender and cause of renal failure

